

## MAMA Project, Inc. Basic MAMA Brigade Kit /100 population

ITEM	Use	#	X
MAMA CHILD SURVIVAL KIT PART A (packed by IMRES)			
Acetaminophen 500mg or Paracetamol tablets (Bags of 30)	1 bag per family	25	
Albendazole 400mg tablets	1 per person	100	
Amoxicillin 250mg tablets	Emergency infection	1000	
Atenolol 25mg	Emergency BP	1000	
Enalapril 10mg	Emergency BP	1000	
Benzyl Benzoate 25% Oil in liter bottles	Scabies-40cc/family	4	
Cimetidine 800mg tablets	Emergency-gastritis	1000	
Ciprofloxacin 500mg	Emergency infection	1000	
Ibuprofen 200mg tablets (Bags of 30)	1 bag per family	25	
Metformin 500mg	Emergency-diabetes	1000	
Metronidazole 250mg tablets	Emergency infection	1000	
Oral rehydration salts envelopes	5 for each family	125	
Prednisone 5mg	Emergency- asthma/severe allergy	200	
Salbutamol 4mg tablets	Emergency asthma	1000	
Sodium Fluoride .25mg	Cavity Prevention	10,000	
Trimethoprim/Sulfamethoxazole 80/400 (Single Strength)	Emergency infection	1000	
Vitamin A 200,000 units/capsule (not for pregnant)	Child survival-as directed	<100	
MAMA CHILD SURVIVAL KIT PART B (packed at MAMA)			
Dentifrice in plastic container 1/family (black oval)	1 container/family	25	
Essential Micronutrient Powder	Bags of 60cc powder	450	
Micronutrient Containers 1/family with scoop and insert	1 container/family	25	
Triple Therapy Cream (Hydrocortisone, Gentamicin, Clotrimazole)	Skin inflammation/infection	25	
Lancets	Anemia test	100	
Alcohol	Anemia test	100	
Blotter paper	Anemia test	100	
Registration sheets 6/100 people	Register Community	6	
Individual record sheet 100	Register each person	100	
Mothers Logs	One for each girl >15	35	
Anemia Color Scale	2 for station	2	
"A Micronutrient Story" DVD (new orders only)	1 for trainer	1	
Pocket Guide for Micronutrient Village Demonstration	1 for trainer	1	
Facilitator's Guide for Micronutrient Village Demonstration	1 for trainer	1	

## MAMA Project, Inc. Basic MAMA Brigade Kit /100 population

ITEM	Use	#	X
ADDENDUM *			
Ergonovine 0.2 mg in a 1mL ampule. 1 or 2 ampules injected into the muscle. Repeat if needed in 30 to 60 minutes than switch to oral when static.	Severe bleeding in delivery		
Ergonovine 0.2mg tablets. To prevent or treat severe bleeding especially if mother is already anemic. Give 1-2 tablets by mouth 4 times daily.			
Oxytocin can be used by injection before delivery of the placenta			
Vitamin K 1 mg per vial (Also called Phytomenadione or phytonadione) To treat bleeding in the newborn or to prevent bleeding in a premature or very small (under 22 kg/ 4.5 lbs) infant, give vitamin K to the baby by injection of one vial only into the baby's outer thigh.			

\*These are a few kinds of medicines that can be life-saving in the hands of a trained midwife. To prevent and treat severe emergency birth complications in mothers and newborns, it is good for the midwife to have a medication to stop severe bleeding after the baby and placenta are delivered.

Rev 2-2013